



Aftercare for Interventional Procedures

- Avoid driving for 2-4 hours
- Let the injection site rest for 2-4 hours
- No heavy lifting for 24 hours
- Normal to have some pain after the local anaesthetic wears off
- Panadol can be used for pain relief if needed
- See your GP if pain persists for more than 24 hours



Aftercare for Interventional Procedures

- Avoid driving for 2-4 hours
- Let the injection site rest for 2-4 hours
- No heavy lifting for 24 hours
- Normal to have some pain after the local anaesthetic wears off
- Panadol can be used for pain relief if needed
- See your GP if pain persists for more than 24 hours



Aftercare for Interventional Procedures

- Avoid driving for 2-4 hours
- Let the injection site rest for 2-4 hours
- No heavy lifting for 24 hours
- Normal to have some pain after the local anaesthetic wears off
- Panadol can be used for pain relief if needed
- See your GP if pain persists for more than 24 hours



Aftercare for Interventional Procedures

- Avoid driving for 2-4 hours
- Let the injection site rest for 2-4 hours
- No heavy lifting for 24 hours
- Normal to have some pain after the local anaesthetic wears off
- Panadol can be used for pain relief if needed
- See your GP if pain persists for more than 24 hours



Aftercare for Interventional Procedures

- Avoid driving for 2-4 hours
- Let the injection site rest for 2-4 hours
- No heavy lifting for 24 hours
- Normal to have some pain after the local anaesthetic wears off
- Panadol can be used for pain relief if needed
- See your GP if pain persists for more than 24 hours



Aftercare for Interventional Procedures

- Avoid driving for 2-4 hours
- Let the injection site rest for 2-4 hours
- No heavy lifting for 24 hours
- Normal to have some pain after the local anaesthetic wears off
- Panadol can be used for pain relief if needed
- See your GP if pain persists for more than 24 hours